

**ANTIBACTERIAL ACTIVITY OF *Caulerpa* sp. AT BLUE  
LAGOON, PORT DICKSON, NEGERI SEMBILAN**

**NOOR AISYAH BINTI ABDULLAH**

**Final Year Project Report Submitted in  
Partial Fulfilment of the Requirements for the  
Degree of Bachelor of Science (Hons.) Biology  
in the Faculty of Applied Sciences  
Universiti Teknologi MARA**

**JANUARY 2014**

This Final Year Project Report entitled “**Antibacterial Activity of *Caulerpa* sp. at Blue Lagoon, Port Dickson, Negeri Sembilan**” was submitted by Noor Aisyah binti Abdullah, in partial fulfilment of the requirements for the Degree of Bachelor of Science (Hons.) Biology, in the Faculty of Applied Sciences, and was approved by

---

Sarini binti Ahmad Wakid  
Supervisor  
Faculty of Applied Sciences  
Universiti Teknologi MARA  
72000 Kuala Pilah  
Negeri Sembilan

---

Sarini binti Ahmad Wakid  
Project Coordinator  
Faculty of Applied Sciences  
Universiti Teknologi MARA  
72000 Kuala Pilah  
Negeri Sembilan

---

Dr. Nor ‘Aishah binti Abu Shah  
Head of Pure Science School  
Faculty of Applied Sciences  
Universiti Teknologi MARA  
72000 Kuala Pilah  
Negeri Sembilan

Date: \_\_\_\_\_

## TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	vi
LIST OF FIGURES	vii
LIST OF ABBREVIATIONS	viii
ABSTRACT	ix
ABSTRAK	x
<b>CHAPTER 1: INTRODUCTION</b>	
1.1 Background of Study	1
1.2 Problem Statement	3
1.3 Significant of Study	3
1.4 Objectives of the Study	4
<b>CHAPTER 2: LITERATURE REVIEW</b>	
2.1 Study Site	5
2.2 Overview of Algae	7
2.2.1 Introduction to Chlorophyta	8
2.3 Disc Diffusion Method	9
2.4 Antibacterial Activity of Algae	10
2.5 Extractions	11
2.5.1 Aqueous extract	11
2.5.2 Ethanol extract	11
2.6 Gram Staining	12
2.7 Bacteria	13
2.7.1 Gram-positive bacteria	13
2.7.1.1 <i>Bacillus subtilis</i>	13
2.7.1.2 <i>Staphylococcus</i> sp.	14
2.7.2 Gram-negative bacteria	15
2.7.2.1 <i>Escherichia coli</i>	15
2.7.2.2 <i>Pseudomonas aeruginosa</i>	16
<b>CHAPTER 3: METHODOLOGY</b>	
3.1 Materials	17
3.1.1 Raw materials	17
3.1.2 Bacteria used	17
3.1.3 Chemicals	17

3.1.4	Apparatus	18
3.2	Methods	19
3.2.1	Sample collection	19
3.2.2	Gram staining	19
3.2.3	Bacterial preparations	20
3.2.4	Sample extractions	21
3.2.4.1	Aqueous extraction	21
3.2.4.2	Ethanol extraction	22
3.2.5	Determination of antibacterial activity of crude extract	23
3.2.5.1	Preparation of nutrient agar	23
3.2.5.2	Disc diffusion assay	23
3.2.6	Measurement of inhibition zones	25
3.3	Statistical Analysis	25

#### **CHAPTER 4: RESULTS AND DISCUSSION**

4.1	Algae Collected and Identification of <i>Caulerpa</i> sp.	27
4.2	Solvent Extractions of <i>Caulerpa racemosa</i>	29
4.3	Antibacteria Activity and Disc Diffusion Assay	31

#### **CHAPTER 5: CONCLUSION AND RECOMMENDATION**

5.1	Conclusions	40
5.2	Recommendations	41

#### **CITED REFERENCES**

42

#### **APPENDICES**

51

#### **CURRICULUM VITAE**

55

## ABSTRACT

### ANTIBACTERIAL ACTIVITY OF *Caulerpa* sp. AT BLUE LAGOON, PORT DICKSON, NEGERI SEMBILAN

The aim of this study is to determine the antibacterial activity of *Caulerpa* sp. collected at Blue Lagoon, Port Dickson, Negeri Sembilan on Gram-positive bacteria (*Bacillus subtilis* and *Staphylococcus* sp.) and Gram-negative bacteria (*Escherichia coli* and *Pseudomonas aeruginosa*). *Caulerpa racemosa* identified at Blue Lagoon was tested for the antibacterial activity against selected bacteria strains. Two types of extraction (aqueous and ethanol) on disc diffusion method was used to evaluate the antibacterial activity of *C. racemosa*. The results of antibacterial activity were expressed as a zone of inhibition (in millimetre) and minimum inhibitory concentrations (MIC). The selected bacteria strains were tested with five different concentrations (2, 5, 10, 15, and 20 mg/ml) of the extracts. All the ethanol and aqueous extracts concentrations of *C. racemosa* did not exhibited any antibacterial activity against all bacteria strains, except against *E. coli* (Gram-negative bacteria) on aqueous extracts. The *E. coli* showed very low sensitivity towards all aqueous extracts concentrations of *C. racemosa*. The statistical analysis of ANOVA showed that both ethanol and aqueous extract did not have significant differences against the bacterial strains used ( $p > 0.05$ ).